

**COMPARISON OF INHIBITORY EFFECTS BETWEEN
NATURAL AND CHEMICAL ANTI-BROWNING
AGENTS ON POLYPHENOL OXIDASE IN APPLE JUICE**

NUR FILZA BINTI MOHD FUAD

**Final Year Project Report Submitted in
Partial Fulfillment of the Requirements for the
Degree of Bachelor of Science (Hons.) Biology
In the Faculty of Applied Sciences
Universiti Teknologi MARA**

JULY 2017

This Final Year Project Report entitled **“Comparison of Inhibitory Effects between Natural and Chemical Anti-browning Agents on Polyphenol Oxidase in Apple Juice”** was submitted by Nur Filza binti Mohd Fuad, in partial fulfilment of the requirements for the Degree of Bachelor of Science (Hons.) Biology, in the Faculty of Applied Sciences, and was approved by

Amirah binti Sharif
Supervisor
B. Sc. (Hons.) Biology
Faculty of Applied Sciences
UiTM Negeri Sembilan
Kampus Kuala Pilah
Pekan Perit Tinggi
72000 Kuala Pilah Negeri Sembilan

Lili Syahani binti Rusli
Project Coordinator
B. Sc. (Hons.) Biology
Faculty of Applied Sciences
UiTM Negeri Sembilan
Kampus Kuala Pilah
Pekan Parit Tinggi
72000 Kuala Pilah
Negeri Sembilan

Dr Nor' aishah binti Abu Shah
Head of Programme
B. Sc. (Hons.) Biology
Faculty of Applied Sciences
UiTM Negeri Sembilan
Kampus Kuala Pilah
Pekan Parit Tinggi
72000 Kuala Pilah
Negeri Sembilan

Date: _____

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF FIGURES	vi
LIST OF ABBREVIATIONS	vii
ABSTRACT	viii
ABSTRAK	ix

CHAPTER 1: INTRODUCTION

1.1	Background of Study	1
1.2	Problem Statement	3
1.3	Significance of the Study	3
1.4	Objectives of the Study	4

CHAPTER 2: LITERATURE REVIEW

2.1	Fruit Juice	5
2.2	Enzymatic Browning	6
2.2.1	Polyphenol oxidase	6
2.3	Phenolic Compounds	8
2.4	Anti-Browning Agents	9
2.4.1	Chemical anti-browning agents	9
2.4.2	Natural anti-browning agents	10

CHAPTER 3: METHODOLOGY

3.1	Materials	12
3.1.1	Raw materials	12
3.1.2	Chemicals	12
3.1.3	Apparatus	12
3.2	Methods	13
3.2.1	Onion extract preparation	13
3.2.2	Polyphenol oxidase enzyme preparation	13
3.2.3	Polyphenol oxidase assay	14
3.2.4	Preparation of gallic acid standard curve	15
3.2.5	Total phenolic content measurement	15
3.3	Data Analysis	16

CHAPTER 4: RESULTS AND DISCUSSION	
4.1 Inhibitory Effect between Ascorbic Acid and Heated Onion Extract on Apple Juice PPO	17
4.2 Determination of Total Phenolic Content in Apple Juice	20
CHAPTER 5: CONCLUSION AND RECOMMENDATIONS	23
CITED REFERENCES	24
APPENDICES	28
CURRICULUM VITAE	31

ABSTRACT

COMPARISON OF INHIBITORY EFFECTS BETWEEN NATURAL AND CHEMICAL ANTI-BROWNING AGENTS ON POLYPHENOL OXIDASE IN APPLE JUICE

Apple juice is known for its good taste and high content of nutrition. However, enzymatic browning easily occurred in it which reduce the product quality and nutrition value caused by activation of polyphenol oxidase. This is due to the damage of cell structure and contact between enzyme and substrate are promoted. Since the chemical anti-browning agent is limited due to short period time function, high cost or cause adverse health effect, development of natural anti-browning agent will be beneficial to overcome the disadvantages. The aim of this study were to compare the inhibitory effect and to determine the total phenolic content in naturally and chemically treated apple juice. The inhibition activity of both anti-browning agents were measured by performing polyphenol oxidase assay on the apple juice. The inhibitory activity of the ascorbic acid on apple juice PPO range from 33.02% to 96.26% compared to heated onion extract treatment which inhibitory activity reached its maximum at 69.16%. The total phenolic content in treated apple juice were determined using the Folin-Ciocalteu method with gallic acid as standard. Phenolic content of apple juice with ascorbic acid treatment was about 9.223 mg GAE/100 ml compared to apple juice treated with heated onion extract was 2.622 mg GAE/100 ml. Ascorbic acid treatment showed higher total phenolic content and lower polyphenol oxidase activity on apple juice compared to heated onion extract. Therefore, it can be concluded that higher total phenol content will cause lower polyphenol oxidase activity.